

Quantum mechanics on moduli spaces.

Ian G. Moss and Noriko Shiiki

Department of Physics, University of Newcastle Upon Tyne, NE1 7RU U.K.

(June 1998)

Abstract

It has been assumed that it is possible to approximate the interactions of quantized BPS solitons by quantising a dynamical system induced on a moduli space of soliton parameters. General properties of the reduction of quantum systems by a Born-Oppenheimer approximation are described here and applied to sigma models and their moduli spaces in order to learn more about this approximation. New terms arise from the reduction procedure, some of them geometrical and some of them dynamical in nature. The results are generalised to supersymmetric sigma models, where most of the extra terms vanish.

Pacs numbers: 03.70.+k, 98.80.Cq

I. INTRODUCTION

In most physical problems, the degrees of freedom that are of interest are only a small fraction of the total number of degrees of freedom present. Some kind of reduction of the system has been performed. The way in which this reduction proceeds from quantum field theory to a moduli space of topological solitons is the subject of this paper.

Soliton solutions can give rise to particle states [1] and it has been predicted that their low-energy interactions can be reduced to a simplified quantum mechanical problem in a set of collective coordinates [2,3]. One way in which this can be used is to indicate a duality between two quantum theories, where the soliton particle states in one theory become the fundamental particles in an equivalent quantum theory. A particular example can be seen in the monopole bound states which are important for the duality between the large and small coupling constant limits of the heterotic string theory [4].

In a range of different models there are classical multiple-soliton solutions that saturate Bogomol'nyi bounds and are parameterised by moduli. When the solitons move slowly and interact they deform adiabatically and the moduli trace out a path in a moduli space. The equations governing the path are equivalent to the equations for a point particle moving on a curved space [5,6].

We would expect to be able to describe the interactions of quantized solitons by quantising the dynamical system induced on the moduli space. We will call this procedure *truncation*. However, the quantisation rules of the truncated theory are often ambiguous without additional assumptions. If we impose general covariance, reflecting the freedom to choose coordinates on the moduli space, then the Hamiltonian operator takes the form

$$-\frac{1}{2}\Delta + \xi R + \dots \quad (1)$$

where Δ is the Laplacian on the moduli space and ambiguities are reduced to the coefficients ξ of the Ricci scalar R and other curvature invariants [7]. These ambiguities can be further reduced by introducing supersymmetry, in which case it has been argued that $\xi = 0$ [8,9].

In this paper we describe the alternative to truncating the full theory, which is to *reduce* it by a Born-Oppenheimer approximation. In a purely bosonic theory we will see that general covariance breaks down in the sense that a potential arises on the moduli space after integrating out the fluctuations that are orthogonal to the moduli space.

In the basic situation we begin with a classical dynamical system in flat space which has a potential V with a set of degenerate minima $\mathcal{M}$. We find that the original quantum theory reduces to a quantum theory on $\mathcal{M}$ with Hamiltonian

$$H_R = -\frac{1}{2}\Delta_{\mathcal{M}} + U + \frac{1}{4}R_{\mathcal{M}} - \frac{1}{8}|\mathbf{k}|^2 + \dots \quad (2)$$

The two purely geometrical terms depend on the intrinsic curvature scalar $R_{\mathcal{M}}$ and the (traced) extrinsic curvature $\mathbf{k}^I$. These geometrical terms were first written down by Maraner [10]. The dynamical terms denoted by U depend upon derivatives of the potential. Those depending on up to two extrinsic derivatives of the potential can be written

$$U = \frac{1}{2}\text{tr}(\omega) + \frac{1}{16}\text{tr}\left(\omega^{-1\parallel}\nabla\omega \cdot \omega^{-1\parallel}\nabla\omega\right) \quad (3)$$

where ω is a matrix, ω^2 is the hessian matrix of V and the covariant derivative is taken tangentially to $\mathcal{M}$.

The application of these results to sigma models is considered in section 3. In these models the original configuration space, consisting of maps from physical space to a curved target space, is both infinite and curved. The infinities can be dealt with by regularisation. The curvature would lead in general to ambiguities even before the reduction. However, as shown in appendix A, these ambiguities are not present if the original space is a Riemannian symmetric space. Most models of interest fall into this class.

The results are generalised to supersymmetric sigma models in section 4. The point at issue is whether the reduction leads to a superpotential on the moduli space and alters the hamiltonian operator. We find that, to leading order, the theory reduces to the Hodge Laplacian acting on p -forms,

$$H_R = -\frac{1}{2}\Delta_{\mathcal{M}} + \dots \quad (4)$$

confirming previous expectations [8,9,3]. Possible developments of this work are addressed in the conclusion, where we also discuss the advantages of the Hamiltonian approach over a Feynmann diagram approach.

II. BORN-OPPENHEIMER REDUCTION

Consider a classical dynamical system in flat space which has a degenerate set of stable points $\mathcal{M}$. If the potential is sufficiently steep, then energy spectrum of the quantum system typically falls into separate bands. An adiabatic approximation scheme can then be used to reduce the quantum system to a quantum system on $\mathcal{M}$.

We shall restrict attention to the case where the potential has an analytic expansion about its minima. We take a set of generalised coordinates (x^a, x^I) such that $x^I = 0$ on $\mathcal{M}$. In the lowest energy bands, the value of x^I fluctuates over a narrow range of values determined by the eigenvalues of a matrix ω , where

$$\omega^2 = \partial_I \partial_J V. \quad (5)$$

The expansion parameter of the approximation scheme is given by the width of an energy band divided by the smallest eigenvalue.

The first step is to rewrite the Hamiltonian operator,

$$H = -\frac{1}{2}\Delta + V \quad (6)$$

in the new coordinate system. We choose normal coordinates x^I along the principal directions of $\partial_I \partial_J V$. It is always possible to find two mutually orthogonal sets of derivatives $\hat{\partial}_a$ and ∂_I , where

$$\hat{\partial}_a = \partial_a - N^I_a \partial_I. \quad (7)$$

This implies that the metric can be written

$$\mathbf{g} = \gamma_{ab} dx^a dx^b + \gamma_{IJ} (dx^I + N^I_a dx^a)(dx^J + N^J_b dx^b). \quad (8)$$

The metric components have normal coordinate expansions

$$\gamma_{IJ} = \delta_{IJ} \quad (9)$$

$$N_a^I = -a_{J^I a} x^J \quad (10)$$

$$\gamma_{ab} = \sigma_{ab} + 2k_{Iab} x^I + k_{Iac} k_{Jb}^c x^I x^J \quad (11)$$

where the extrinsic curvature $\mathbf{k}^I$ and the torsion $\mathbf{a}_I^J$ are defined in appendix B. These expansions are exact if the space is flat, but we will keep in mind the possibility of generalising these results to curved spaces.

In the new coordinate system, the Hamiltonian becomes

$$H = \frac{1}{2}|g|^{-1/2} \hat{\partial}_a |g|^{1/2} \gamma^{ab} \hat{\partial}_b + \frac{1}{2}|g|^{-1/2} \partial_I |g|^{1/2} \delta^{IJ} \partial_J + V \quad (12)$$

We regard the x^I as small and expand the Hamiltonian as a series of terms $H = H_0 + H_1 + \dots$. The potential has a series expansion

$$V = \frac{1}{2}(\omega^2)_{IJ} x^I x^J + \frac{1}{6} V_{IJK} x^I x^J x^K + \dots \quad (13)$$

where ω_{IJ} is a diagonal matrix with eigenvalues ω_I . The first few terms in the full Hamiltonian are then

$$H_0 = -\frac{1}{2} \delta^{IJ} \partial_I \partial_J + \frac{1}{2} (\omega^2)_{IJ} x^I x^J \quad (14)$$

$$H_1 = -\frac{1}{2} \text{tr}(\mathbf{k}^I) \partial_I + \frac{1}{6} V_{IJK} x^I x^J x^K \quad (15)$$

$$H_2 = -\frac{1}{2} \mathcal{D} \cdot \mathcal{D} + \frac{1}{2} \text{tr}(\mathbf{k}_I \mathbf{k}^J) x^I \partial_J + \frac{1}{24} V_{IJKL} x^I x^J x^K x^L \quad (16)$$

The dot denotes a scalar product using the intrinsic metric on $\mathcal{M}$. We have introduced a covariant derivative $\mathcal{D}_a$, where $\mathcal{D}_a \sigma_{ab} = 0$ and

$$\mathcal{D}_a f = (\partial_a + a_{J^I a} x^J \partial_J) f \quad (17)$$

for scalar functions f .

The next step is to apply degenerate perturbation theory. We take the unperturbed harmonic oscillator Hamiltonian H_0 together with a set of states f_n , where $H_0 f_n = E_n f_n$. The perturbation theory is described in appendix C and gives a reduced Hamiltonian operator H_R acting on the space of wave functions $\psi(x^a)$,

$$H_R = \langle H_0 \rangle_{00} + \langle H_1 \rangle_{00} + \langle H_2 \rangle_{00} + \sum_{m \neq 0} \langle H_1 \rangle_{0m} (E_0 - E_m)^{-1} \langle H_1 \rangle_{m0} + O(\omega^{-1}) \quad (18)$$

The matrix elements defined by

$$\langle H \rangle_{nm} = \int f_n^* H f_m \prod_I dx^I \quad (19)$$

are allowed to be operator-valued.

The first term $\langle H_0 \rangle_{00}$ consists of the vacuum energy of the system of unperturbed harmonic oscillators (14),

$$E_0 = \frac{1}{2} \text{tr}(\omega). \quad (20)$$

The second term $\langle H_1 \rangle_{00}$ vanishes because the operator is odd under the inversion $x^I \rightarrow -x^I$. For the next term $\langle H_2 \rangle_{00}$, with H_2 given by equation (16), we set

$$\langle \mathcal{D}_a \rangle_{mn} = \delta_{mn} \parallel \nabla_a + \langle \mathcal{A}_a \rangle_{mn} \quad (21)$$

where $\parallel \nabla_a$ is defined in appendix B and $\mathcal{A}$ arises from the action of $\mathcal{D}_a$ on the states f_n in equation (19). The matrix element of the derivative of an energy eigenstate is given by a standard identity,

$$\langle \mathcal{A}_a \rangle_{mn} = \frac{([\mathcal{D}_a, H_0])_{mn}}{E_n - E_m} \quad (22)$$

After using equation (17) for $\mathcal{D}$, and comparing with equation (B7), the commutator produces a covariant derivative,

$$\langle \mathcal{A}_a \rangle_{mn} = \begin{cases} \frac{1}{2} \parallel \nabla_a \omega_{IJ} \langle x^I x^J \rangle_{mn} & E_m < E_n \\ 0 & m = n \end{cases} \quad (23)$$

We can now evaluate

$$\langle \mathcal{D} \cdot \mathcal{D} \rangle_{00} = \Delta_{\mathcal{M}} - \frac{1}{8} \text{tr} \left(\omega^{-1} \parallel \nabla \omega \cdot \omega^{-1} \parallel \nabla \omega \right) \quad (24)$$

where the covariant Laplacian $\Delta_{\mathcal{M}} = \parallel \nabla \cdot \parallel \nabla$. We also have, using (B8),

$$\langle \text{tr}(\mathbf{k}_I \mathbf{k}^J) x^I \partial_J \rangle_{00} = \frac{1}{2} \text{tr}(\mathbf{k}_I \mathbf{k}^I) = \frac{1}{2} |\mathbf{k}|^2 - \frac{1}{2} R_{\mathcal{M}} \quad (25)$$

where $|\mathbf{k}|^2 = \text{tr}(\mathbf{k}^I) \text{tr}(\mathbf{k}_I)$. Therefore we get

$$\langle H_2 \rangle_{00} = -\frac{1}{2} \Delta_{\mathcal{M}} - \frac{1}{4} |\mathbf{k}|^2 + \frac{1}{4} R_{\mathcal{M}} + \frac{1}{16} \text{tr} \left(\omega^{-1} \parallel \nabla \omega \cdot \omega^{-1} \parallel \nabla \omega \right) + \frac{1}{32} \sum_{IJ} \omega_I^{-1} \omega_J^{-1} V_{IJIJ} \quad (26)$$

The last term given in equation (18) leads to

$$\frac{1}{8} |\mathbf{k}|^2 - \frac{1}{48} \sum_{IJK} (\omega_I + \omega_J + \omega_K)^{-1} \omega_I^{-1} \omega_J^{-1} \omega_K^{-1} (V_{IJK})^2 - \frac{1}{32} \sum_{IJK} \omega_I^{-2} \omega_J^{-1} \omega_K^{-1} V_{IJJ} V_{IKK} \quad (27)$$

Putting the terms together gives a final expression for the reduced Hamiltonian,

$$H_R = -\frac{1}{2} \Delta_{\mathcal{M}} + U + \frac{1}{4} R_{\mathcal{M}} - \frac{1}{8} |\mathbf{k}|^2 + O(\omega^{-1}) \quad (28)$$

The terms are divided into purely geometrical terms and dynamical terms which are collected into U , where

$$U = \frac{1}{2} \text{tr}(\omega) + \frac{1}{16} \text{tr} \left(\omega^{-1} \parallel \nabla \omega \cdot \omega^{-1} \parallel \nabla \omega \right) + \frac{1}{32} \sum_{IJ} \omega_I^{-1} \omega_J^{-1} V_{IJIJ} \\ - \frac{1}{48} \sum_{IJK} (\omega_I + \omega_J + \omega_K)^{-1} \omega_I^{-1} \omega_J^{-1} \omega_K^{-1} V_{IJK}^2 - \frac{1}{32} \sum_{IJK} \omega_I^{-2} \omega_J^{-1} \omega_K^{-1} V_{IJJ} V_{IKK} \quad (29)$$

The vacuum energy, which is of order ω , is the dominant term in this expression. The next term depends on the gradient of the normal mode frequencies along the moduli space and includes the effects of the twisting, or torsion, of the normal mode directions. The remaining terms contain the effects of higher derivatives of the potential.

The geometrical terms were discovered in the restricted situation of a surface embedded in R^3 by Jensen and Koppe [11] and more generally by Maraner [10]. The covariant forms for the dynamical terms beyond the obvious vacuum energy term are new, as far as we know.

III. BOSONIC FIELD THEORY

The non-linear sigma-model in two space dimension x^μ and one time dimension t has fields ϕ^i taking values in a curved target space $\mathcal{T}$. We will take the Lagrangian to be $L = T - V$, where

$$T = \frac{1}{2} \int g_{ij} \partial_t \phi^i \partial_t \phi^j d^2x \quad (30)$$

$$V = \frac{1}{2} \int g_{ij} \partial_\mu \phi^i \partial_\mu \phi^j d^2x \quad (31)$$

Depending upon the topology of $\mathcal{T}$, the classical field equations can have static solutions with finite energy, or topological solitons. These solutions minimise the potential energy for a given topological class.

When $\mathcal{T}$ is a compact Kahler manifold the soliton solutions can be represented by holomorphic maps from the complex plane into $\mathcal{T}$ [6,12,13]. They depend on a continuous set of parameters, or moduli, which can be interpreted as the positions and charges of individual lumps.

The initial situation is similar to the models considered in the previous section, except that the configuration space is no longer finite and flat. The introduction of curvature leads to the appearance of curvature invariants in the initial Schrödinger equation. However, we will take the target space to be a Riemannian symmetric space. The space of fields inherits this symmetry and is also a Riemannian symmetric space. In this case the curvature invariants are constants which can be absorbed into an overall phase factor.

The soliton solutions ϕ_0 will be parameterised by a set of coordinates x^a belonging to the moduli space. We shall expand the Lagrangian about these solutions using standard background field methods [14] and express the result in terms of time-dependent coordinates x^a and normal coordinates x^I . This will enable us to use the results of the previous section.

The tangent vector ξ to the geodesic from ϕ to ϕ_0 provides a convenient measure of the displaced field. We also introduce the vectors $\mathbf{e}_\mu$ with target space components

$$\mathbf{e}_\mu{}^i = \partial_\mu \phi^i \quad (32)$$

These can be parallel transported back from the tangent space at ϕ to the tangent space at ϕ_0 to produce a field $\mathbf{e}'_\mu$. The tangent vector ξ commutes with $\mathbf{e}_\mu$, and consequently $\nabla_\xi \mathbf{e}_\mu = D_\mu \xi$, where D_μ is the gradient along $\mathbf{e}_\mu$. This relation simplifies the Taylor series expansion for $\mathbf{e}'$,

$$\mathbf{e}'_\mu = \mathbf{e}_\mu + D_\mu \xi + \frac{1}{2} R(\xi, \mathbf{e}_\mu) \xi + \frac{1}{6} R(\xi, D_\mu \xi) \xi + \dots \quad (33)$$

We will define

$$(\mathbf{u}, \mathbf{v}) = \int g_{ij}(\phi_0) u^i v^j d^2x \quad (34)$$

By parallel transport of the Lagrangian density we can write the potential in the form

$$V = \frac{1}{2} (\mathbf{e}'_\mu, \mathbf{e}'_\mu) \quad (35)$$

The potential therefore has a series expansion in ξ of the form $V = V_0 + V_1 + V_2 \dots$ where

$$V_0 = \frac{1}{2}(\mathbf{e}_\mu, \mathbf{e}_\mu) \quad (36)$$

$$V_1 = (\mathbf{e}_\mu, D_\mu \xi) \quad (37)$$

$$V_2 = \frac{1}{2}(D_\mu \xi, D_\mu \xi) + \frac{1}{2}(\mathbf{e}_\mu, R(\xi, \mathbf{e}_\mu)\xi) \quad (38)$$

$$V_3 = \frac{2}{3}(D_\mu \xi, R(\xi, \mathbf{e}_\mu)\xi) \quad (39)$$

$$V_4 = \frac{1}{6}(D_\mu \xi, R(\xi, D_\mu \xi)\xi) + \frac{1}{6}(R(\xi, \mathbf{e}_\mu)\xi, R(\xi, \mathbf{e}_\mu)\xi) \quad (40)$$

Integration by parts in (37) produces the field equation

$$D_\mu \mathbf{e}_\mu = \partial_\mu e_\mu^i - \Gamma_{jk}^i e_\mu^j e_\mu^k = 0 \quad (41)$$

Integration by parts on (38) produces the operator

$$\Delta_f \xi = -D_\mu D_\mu \xi - R(\xi, \mathbf{e}_\mu)\mathbf{e}_\mu, \quad (42)$$

which describes fluctuations about the moduli space. In the usual analysis of the sigma model, the inverse of this propagator would be the Green function in the background field approximation.

The normalised positive modes of the fluctuation operator will be denoted by $\mathbf{u}_I$ and their eigenvalues by $(\omega_I)^2$. The fluctuation operator also has a set of zero modes $\mathbf{u}_a = \partial_a \phi_0$. All of the modes are parameterised by the collective coordinates x^a . We define the remaining coordinates in terms of the displacement vector,

$$\xi = x^I \mathbf{u}_I \quad (43)$$

The potential then has a series expansion in the normal coordinates identical to equation (13) used in the previous section. From equation (39),

$$V_{IJK} = 4C_{(IJ)K} \quad (44)$$

where

$$C_{IJK} = (D_\mu \mathbf{u}_I, R(\mathbf{u}_J, \mathbf{e}_\mu)\mathbf{u}_K). \quad (45)$$

The coefficients are symmetrical, since $C_{(IJ)K} = C_{(IJK)}$. They are also trace-free if the zero modes u_a are included, $C^I_{IJ} + C^a_{aJ} = 0$.

The kinetic energy has a similar series expansion. Since

$$T = \frac{1}{2}(\mathbf{e}'_t, \mathbf{e}'_t), \quad (46)$$

we need only replace $\mathbf{e}_\mu$ by $\mathbf{e}_t$ and $D_\mu \xi$ by $D_t \xi$ in equations (36-39). However, both the x^I and the x^a depend on time, so that the time derivative of equation (43) implies

$$D_t \xi = \dot{x}^I \mathbf{u}_I + \dot{x}^a x^I D_a \mathbf{u}_I \quad (47)$$

where the covariant derivative

$$D_a u^i = \partial_a u^i + u_a^k \Gamma_{jk}^i u^j. \quad (48)$$

Also, since $\mathbf{u}_a = \partial_a \phi_0$,

$$\mathbf{e}_t = \partial_t \phi_0 = \dot{x}^a \mathbf{u}_a. \quad (49)$$

Equations (47) and (49) allow terms in the series expansion of the kinetic energy to be grouped as a quadratic polynomial in the generalised velocities,

$$T = \frac{1}{2} \gamma_{ab} \dot{x}^a \dot{x}^b + \frac{1}{2} \gamma_{IJ} (\dot{x}^I + N^I_a \dot{x}^a) (\dot{x}^J + N^J_b \dot{x}^b) \quad (50)$$

To second order in x^I ,

$$\gamma_{IJ} = \delta_{IJ} - \frac{1}{3} R_{IKJL} x^K x^L \quad (51)$$

$$N^I_a = -a_J^I x^J \quad (52)$$

$$\gamma_{ab} = \sigma_{ab} + 2k_{Iab} x^I + k_{Iac} k_{Jb}^c x^I x^J - R_{aIbJ} x^I x^J \quad (53)$$

The coefficients can be obtained by examining equations (36-39),

$$\sigma_{ab} = (\mathbf{u}_a, \mathbf{u}_b) \quad (54)$$

$$k_{ab}^I = (\mathbf{u}_a, D_b \mathbf{u}_I) \quad (55)$$

$$a_I^J{}_a = (\mathbf{u}_I, D_a \mathbf{u}_J) \quad (56)$$

$$R_{IJKL} = (\mathbf{u}_I, R(\mathbf{u}_K, \mathbf{u}_L) \mathbf{u}_J) \quad (57)$$

These results recover the normal coordinate expansion of the previous section (9)-(11), with extra terms due to the curvature of the original configuration space. We now have explicit expressions for the curvature and the extrinsic geometry of the space of soliton parameters in terms of eigenstates of the fluctuation operator.

The results of the previous section still apply, except for the addition of terms depending on the target space curvature. A similar analysis to the previous section shows that two extra terms

$$\frac{1}{4} \sigma^{ac} \sigma^{bd} R_{abcd} + \frac{1}{6} \sum_{IJ} \omega_I \omega_J^{-1} R_{IJIJ} \quad (58)$$

should be included in the potential U .

Expressions involving derivatives of the eigenmodes can be rewritten using the identity

$$(\mathbf{u}_I, D_a \mathbf{u}_J) = \frac{(\mathbf{u}_I, [D_a, \Delta_f] \mathbf{u}_J)}{\omega_J^2 - \omega_I^2} \quad (59)$$

With the fluctuation operator Δ_f given explicitly in equation (42), the commutator becomes

$$(\mathbf{u}_I, [D_a, \Delta_f] \mathbf{u}_J) = 4C_{(IJ)a} \quad (60)$$

where C_{IJa} is defined as in equation (45). This gives

$$k_{Iab} = -\frac{4}{\omega_I^2} C_{(Ia)b} \quad (61)$$

$$\|\nabla_a \omega_{IJ} = \frac{4}{\omega_J + \omega_I} C_{(IJ)a} \quad (62)$$

where equation (B7) has been used.

The dominant term in the reduced Hamiltonian should be the zero point energy $\frac{1}{2} \text{tr}(\omega)$. In general, in order to evaluate this term it will be necessary to solve the eigenvalue problem for the fluctuation operator numerically. Once this has been done, equations (44), (61) and (62) could then be used to evaluate the geometrical terms in the reduced Hamiltonian (28).

IV. SUPERSYMMETRIC REDUCTION

We now turn to the reduction of an $N = 2$ supersymmetric quantum system defined on a Riemannian symmetric space. As before, we suppose that the system has a potential V with a degenerate set of minima. The existence of two supersymmetries requires that the space is a Kahler manifold with a covariantly preserved complex structure J [15,16]. We shall suppose that the moduli space of stable points is also a Kahler manifold.

The variables consist of coordinates x^α and single complex component fermions ψ^α . We also assume the existence of an operator π_α , related to the momentum, and the following commutation relations

$$\{\psi^\alpha, \psi^{*\beta}\} = g^{\alpha\beta} \quad (63)$$

$$[\pi_\alpha, x^\beta] = -i\delta_\alpha^\beta \quad (64)$$

$$[\pi_\alpha, \psi^\beta] = -i\Gamma^\beta_{\gamma\alpha}\psi^\gamma \quad (65)$$

$$[\pi_\alpha, \pi_\beta] = -iR_{\alpha\beta\gamma\delta}\psi^{*\gamma}\psi^\delta \quad (66)$$

involving the inverse metric $g^{\alpha\beta}$, connection components $\Gamma^\beta_{\gamma\alpha}$ and curvature components $R_{\alpha\beta\gamma\delta}$.

It is possible to represent the commutator relations by covariant differential operators on a Hilbert space with the basis

$$\Psi(x)_{\alpha_1 \dots \alpha_p} \psi^{*\alpha_1} \dots \psi^{*\alpha_p} |0\rangle, \quad (67)$$

where $\psi^\alpha |0\rangle = 0$. This gives a representation in which

$$\pi_\alpha = -i(\partial_\alpha - \Gamma_{\gamma\beta\alpha}\psi^{*\beta}\psi^\gamma) \quad (68)$$

The action of π is equivalent to $-i\nabla_\alpha$ acting on the p -forms Ψ , where ∇_α is the metric connection.

We shall assume that the classical system has two supersymmetries generated by the supercharges

$$Q^{(1)} = \psi^\alpha \pi_\alpha - \psi^{*\alpha} W_\alpha(x) \quad (69)$$

$$Q^{(2)} = (J\psi)^\alpha \pi_\alpha - (J\psi^*)^\alpha W_\alpha(x) \quad (70)$$

The reason for choosing this particular form for these supercharges will become apparent when we come to discuss sigma models in the next section. We will only require the combination

$$Q = \frac{1}{2}(Q^{(1)} + iQ^{(2)}) = \lambda^\alpha \pi_\alpha - \nu^\alpha W_\alpha(x) \quad (71)$$

where the fermion fields

$$\lambda = \frac{1}{2}(1 + iJ)\psi \quad (72)$$

$$\nu = \frac{1}{2}(1 + iJ)\psi^* \quad (73)$$

are purely holomorphic.

The Hamiltonian is given (up to a constant) by $H = \{Q, Q^*\}$, which can be evaluated using the commutation relations (66),

$$H = -\frac{1}{2}\nabla^2 + g^{\alpha\beta}W_\alpha W_\beta^* + iW_{\alpha;\beta}^*\lambda^\beta\nu^{*\alpha} + iW_{\alpha;\beta}\lambda^{*\beta}\nu^\alpha - \frac{1}{4}R_{\alpha\beta\gamma\delta}[\lambda^\alpha, \lambda^{*\beta}][\lambda^{*\gamma}, \lambda^\delta] - \frac{1}{4}R_{\alpha\beta\gamma\delta}[\lambda^\alpha, \lambda^{*\beta}][\nu^\gamma, \nu^{*\delta}] \quad (74)$$

The moduli space lies at the minimum of the bosonic potential and is parameterised by a set of complex coordinates z^a . We assume that the superpotential W_α has a series expansion in normal coordinates z^I ,

$$W_\alpha = W_{\alpha\bar{J}}z^{\bar{J}} + W_{\alpha\bar{J}\bar{K}}z^{\bar{J}}z^{\bar{K}} + W_{\alpha\bar{J}K\bar{L}}z^{\bar{J}}z^Kz^{\bar{L}} + \dots \quad (75)$$

The normal coordinates can be chosen to ensure that the boson mass matrix is diagonal,

$$g^{\alpha\beta}W_{\alpha\bar{J}}W_{\beta K}^* = (\omega^2)_{\bar{J}K} \quad (76)$$

For the fermion mass matrix $W_{\alpha;\beta}$, we need two separate sets of basis vectors, in general, in order to put it into diagonal form. We can always choose one basis ($\mathbf{e}_a, \mathbf{e}_I$) to coincide with the coordinate basis at $z^I = 0$ for the fields λ . Another basis ($\mathbf{e}_{a'}, \mathbf{e}_{I'}$) then has to be used for the fields ν . (This procedure is similar to the independent unitary rotations of the left and right chirality fermions in the standard model). With the choice

$$\mathbf{e}_{I'} = \omega_I^{-1}W^\alpha_I\mathbf{e}_\alpha, \quad (77)$$

the fermion mass matrix becomes diagonal,

$$W_{I'\bar{J}} = \omega_{I\bar{J}}. \quad (78)$$

The fermion fields λ^a and $\nu^{a'}$ are massless and remain in the reduced quantum system. However, there is an important difference between the finite and infinite dimensional situations. In the infinite dimensional situation the basis $\mathbf{e}_{I'}$ may be complete (in which case there are no massless $\nu^{a'}$ fermions). We shall proceed assuming this to be the case.

Previously, we used a series expansion for the Hamiltonian to arrive at the reduced theory. However, as shown in appendix C, for a supersymmetric theory we need only consider the supercharge. The supercharge (71) depends on the normal coordinates through the superpotential and through the connection coefficients in π_α . In the basis $\mathbf{e}_\alpha = (\mathbf{e}_a, \mathbf{e}_I)$,

$$\Gamma_{\alpha\beta a}\psi^{*\alpha}\psi^\beta \sim \Gamma_{b\bar{c}a}\lambda^{\bar{c}}\lambda^b + k_{\bar{I}ab}\lambda^{\bar{I}}\lambda^a + a_{I\bar{J}a}\lambda^{\bar{J}}\lambda^I + a'_{\bar{I}'J'a}\nu^{J'}\nu^{\bar{I}'} \quad (79)$$

in the limit $z^I = 0$.

The first few terms in the series expansion of the supercharge (71) are now

$$Q_0 = -i\lambda^I\partial_I - \nu^{I'}W_{I'\bar{J}}z^{\bar{J}} \quad (80)$$

$$Q_1 = -i\lambda^a\mathcal{D}_a - \nu^{I'}W_{I'\bar{J}\bar{K}}z^{\bar{J}}z^{\bar{K}} \quad (81)$$

$$Q_2 = -i\lambda^\alpha R_{\beta\gamma\bar{I}\alpha}z^{\bar{I}}(\lambda^{*\gamma}\lambda^\beta - \nu^\gamma\nu^{*\beta}) + \nu^{I'}W_{I'\bar{J}\bar{K}\bar{L}}z^{\bar{J}}z^{\bar{K}}z^{\bar{L}} \quad (82)$$

where $\mathcal{D}_a$ now includes the connection terms (79).

The full quantum theory can now be reduced to the moduli space by following the methods described in appendix C. We define the fermion vacuum state $|0_T\rangle$ which is annihilated by two sets of annihilation operators

$$\lambda_-^I = \frac{1}{\sqrt{2}}(\lambda^I + i\nu^{I'}), \quad \lambda_-^{\bar{I}} = \frac{1}{\sqrt{2}}(\lambda^{\bar{I}} + i\nu^{\bar{I}'}) \quad (83)$$

As in section 2, we build up from $|0_T\rangle$ a set of oscillator states f_n which satisfy $H_0 f_n = E_n f_n$, where H_0 is the unperturbed Hamiltonian. In the supersymmetric case, $E_0 = 0$.

The reduced Hamiltonian is given in terms of the reduced supercharge by

$$H_R = \{Q_R, Q_R^*\} \quad (84)$$

where

$$Q_R = \langle Q_1 \rangle_{00} + \sum_{n \neq 0} \langle Q_1 \rangle_{0n} (E_0 - E_n)^{-1} \langle H_1 \rangle_{n0} + O(\omega^{-1}) \quad (85)$$

The covariant derivatives are treated as in section 2, that is we write

$$\langle \mathcal{D}_a \rangle_{0n} = \delta_{0n} \parallel \nabla_a + \langle \mathcal{A}_a \rangle_{0n} \quad (86)$$

In the supersymmetric case we include the fermion terms from equation (79),

$$\langle \mathcal{A}_a \rangle_{0n} = \parallel \nabla_a \omega_{\bar{I}J} \langle z^{\bar{I}} z^J \rangle_{0n} - k_{\bar{I}ba} \langle \lambda^{\bar{I}} \lambda^b \rangle_{0n} - a_{I\bar{J}a} \langle \lambda^{\bar{J}} \lambda^I \rangle_{0n} + a'_{I'\bar{J}'a} \langle \nu^{I'} \nu^{\bar{J}'} \rangle_{0n} \quad (87)$$

At leading order we recover the same result that we would obtain by a trivial truncation of the theory,

$$Q_R = -i\lambda^a \parallel \nabla_a + O(\omega^{-1}) \quad (88)$$

As discussed in appendix A, the reduced states can be identified with antiholomorphic forms on the moduli space and the reduced Hamiltonian can be identified with the Laplacian. At this order the reduced theory is identical with the truncated theory obtained under the simplest assumptions [3].

V. SUPERSYMMETRIC SIGMA MODELS

The supersymmetric sigma model in two space and one time dimension has Lagrangian

$$L = \frac{1}{2} \int \left(-g_{ij} \partial^\mu \phi^i \partial_\mu \phi^j + i g_{ij} \bar{\chi}^i \gamma^\mu D_\mu \chi^j \right) d^2x \quad (89)$$

where $\mu = 0, 1, 2$, and χ is a two-component majorana spinor. If the target space is a Kähler manifold, with complex structure J , then the model has two supersymmetries. We shall consider the reduction of this theory.

It is convenient to use a complex representation for the fermion fields, with

$$\chi = \begin{pmatrix} \psi \\ \psi^* \end{pmatrix}, \quad (90)$$

and a complex spatial coordinate $z = x + iy$. There are two complex supercharges in this representation,

$$Q^{(1)} = \int \left(g_{ij} \psi^i \partial_t \phi^j - 2g_{ij} \psi^{*i} \partial_z \phi^j \right) d^2 z \quad (91)$$

$$Q^{(2)} = \int \left(g_{ij} (J\psi)^i \partial_t \phi^j - 2g_{ij} (J\psi)^{*i} \partial_z \phi^j \right) d^2 z \quad (92)$$

These can be used to generate the classical supersymmetry transformations by transforming to phase space and using Dirac brackets. The supercharges produce the Hamiltonian by Dirac brackets

$$H = \frac{i}{2} \{Q^{(1)}, Q^{(1)*}\}_{\text{DB}} = \frac{i}{2} \{Q^{(2)}, Q^{(2)*}\}_{\text{DB}} \quad (93)$$

The combination $Q = (Q^{(1)} + iQ^{(2)})/2$ can also be used, but in this case

$$i\{Q, Q^*\}_{\text{DB}} = H + E_T \quad (94)$$

where

$$E_T = 2 \int g_{ij} (J\partial_z \phi)^i (\partial_z \phi)^j d^2 z. \quad (95)$$

The integral is a constant for fields ϕ in the same topological class. We can still use (94) to generate the quantum Hamiltonian.

The background field expansion used in section 3 can be used again here to expand the supercharge Q . The bosonic fluctuations about the background ϕ_0 are described by a vector ξ and a fluctuation operator Δ_f . The fermion fields are simply parallel propagated to ϕ_0 and decomposed into $\psi = \lambda + \nu^*$, where λ and ν are holomorphic, i.e. $J\lambda = i\lambda$.

Since the complex structure J commutes with the fluctuation operator, we can choose holomorphic eigenmodes u_α and use these as a basis $\mathbf{e}_\alpha$. The supercharge for this theory is given by

$$Q = \int (\lambda^i \pi_i - 2g_{ij} \nu^i \partial_z \phi^j) d^2 z \quad (96)$$

When the eigenmode expansions are used, we recover the expression (71) used in section 4,

$$Q = \lambda^\alpha \pi_\alpha - \nu^\alpha W_\alpha(\xi) \quad (97)$$

However, now we have an explicit formula for the superpotential

$$W_\alpha = 2(u_\alpha, \exp(\nabla_\xi) \mathbf{e}_z) \quad (98)$$

in terms of the product (34). The series expansion for W_α in the coordinate z^I is obtained using equation (33). The linear term is,

$$W_{\alpha\bar{J}} = 2(u_\alpha, D_z u_J^*). \quad (99)$$

where

$$D_z u_I = \partial_z u_I^i + (\partial_z \phi_0^j) \Gamma_{jk}^i u_I^k \quad (100)$$

According to equation (76), this implies that the fluctuation operator is

$$\Delta_f u_I = -4D_z D_{\bar{z}} u_I \quad (101)$$

and also that the fermion mass matrix is diagonalised by choosing a basis

$$\mathbf{e}_{I'} = 2\omega_I^{-1} D_{\bar{z}} u_I. \quad (102)$$

These vectors actually form a complete basis, that is they form a normalisable set of eigenvectors of the (positive) operator $D_{\bar{z}} D_z$. The only massless fermions are the λ^a and these remain as fermions on the moduli space. This is precisely the situation considered in the previous section. The reduced theory is therefore identical to the truncated theory at leading order.

VI. CONCLUSION

The reduction of a classical dynamical system generally involves eliminating the internal forces and introducing generalised coordinates. In quantum theory, the uncertainty principle implies that the internal coordinates can never be frozen, but the reduction can be performed approximately when the internal degrees of freedom remain close to their ground state.

We have considered the adiabatic reduction of various quantum systems onto a moduli space of collective coordinates. The results for a quantum mechanical system are quite general and depend on geometrical properties of the moduli space as well as the frequencies of the internal modes. The reduced Hamiltonian to leading order can be found at the end of section 1.

The moduli space of sigma-model solitons has been quantised by taking the continuum limit of the quantum mechanical system. In the Bosonic case, the terms in the reduced Hamiltonian operator depend on the eigenfunctions of a fluctuation operator Δ_f . The spectrum of the fluctuation operator is continuous and the eigenvalue sums of the quantum mechanical system have to be replaced by integrals and regularised.

We could have attempted to quantise the sigma-model moduli space with the instanton techniques used in quantum field theory. In this approach, based on a path integral, the field is replaced by collective coordinates x^a and field fluctuations ξ . The fluctuations are integrated out, leaving a path integral over the coordinates x^a and a Jacobian factor. The classical action S gets replaced by

$$W[x^a] = S[x^a] + \frac{1}{2} \log \det(\Delta_f - \partial_t^2)' + \dots \quad (103)$$

where the prime indicates omission of any zero eigenvalues and the dots denote contributions from higher loop Feynman diagrams. An adiabatic reduction can now be obtained by taking the time derivatives to be small and evaluating some of the higher loop terms. In practice, we have found this approach less practical and not as rigorous as the Hamiltonian one presented earlier.

The bosonic sigma model considered in section 3 can be generalised in various ways. An important possibility would be to include potentials of the form

$$\Phi = \int W(\phi) d^2x. \quad (104)$$

The changes to the results brought about by introducing potentials are confined to the terms in equation (28) involving V_{IJK} and V_{IJKL} , which depend on functional derivatives of Φ .

The supersymmetric sigma model is tightly constrained and reduces to the Hodge Laplacian acting on antiholomorphic p -forms, confirming a result that has been used previously with strong support but not derived rigourously [8,9,3]. It is here, especially, that higher order terms in the adiabatic expansion might be of interest.

APPENDIX A: FACTOR ORDERING

When constructing a quantum theory the ordering of operators is an important issue. A minimum requirement for choosing the operator ordering is to try to retain as much of the symmetry of the classical system as possible. We shall consider symmetry under general coordinate transformations, supersymmetry and the symmetry of a Riemannian Symmetric space.

Our first example is classical particle mechanics in curved space, described by the Hamiltonian

$$H = \frac{1}{2} g^{\alpha\beta} p_\alpha p_\beta. \quad (A1)$$

If we introduce a Hilbert space of wave functions $\psi(x)$ and use the metric to construct an inner product on the Hilbert space, then the Hamiltonian operator

$$H = -\frac{1}{2} |g|^{-1/2} \partial_\alpha |g|^{1/2} g^{\alpha\beta} \partial_\beta. \quad (A2)$$

is both covariant and consistent with the basic principals of quantum mechanics. However, many other possibilities exist, since we are free to add terms such as

$$i\xi R^\alpha{}_\beta [x^\alpha, p_\beta] \quad (A3)$$

which are of order $\hbar$ or higher and vanish in the classical limit. If we assume that the Hamiltonian operator is a covariant second order operator, then the form of this operator must be [7]

$$H = -\frac{1}{2} (\nabla + A)^2 + X \quad (A4)$$

However, the vector field A and scalar field X cannot be inferred from the classical Hamiltonian.

The quantum theory is far more restrictive if the space is a Riemannian Symmetric space. A Riemannian Symmetric space is defined by its geodesic symmetries. The geodesic symmetries about a point leave the point unchanged and map each of the geodesics through the point into itself. A Riemannian Symmetric space has these symmetries at every point and covariant derivatives of the curvature, which are odd under the reflection symmetry, must vanish [19]. In this case, both ∇X and A , which are also odd under the geodesic symmetry, must vanish. The Hamiltonian on a Riemannian symmetric space is therefore given by the Laplacian and a constant.

Our second example is the supersymmetric particle mechanics on a Kahler manifold, originally discussed by Witten [8]. The coordinates z^a and momenta π_a are supplemented by a set of complex fermions λ^a . The commutation relations are

$$\{\lambda^a, \lambda^{\bar{b}}\} = g^{a\bar{b}} \quad (\text{A5})$$

$$[\pi_a, z^b] = -i\delta_a^b \quad (\text{A6})$$

$$[\pi_a, \lambda^b] = i\Gamma_{ca}^b \lambda^c \quad (\text{A7})$$

$$[\pi_a, \pi_{\bar{b}}] = iR_{a\bar{b}d\bar{c}} \lambda^{\bar{c}} \lambda^d \quad (\text{A8})$$

and their hermitian conjugates, where $g^{a\bar{b}}$ is the inverse metric, $\Gamma_{ca}^b \lambda^c$ the connection components and $R_{a\bar{b}d\bar{c}}$ the Riemannian curvature. The supersymmetry is generated by a supercharge

$$Q = \lambda^a \pi_a. \quad (\text{A9})$$

The ordering of the operators in the last commutation relation and the supercharge have so far been chosen arbitrarily.

The commutation relations can be represented by differential operators acting on the Hilbert space of functions $\Psi(z^a, z^{\bar{a}}, \lambda^{\bar{a}})$, with

$$\pi_a = -i\partial_a \quad (\text{A10})$$

$$\pi_{\bar{a}} = -i(\partial_{\bar{a}} - \Gamma_{c\bar{a}}^{\bar{b}} \lambda^{\bar{b}} \lambda^c) \quad (\text{A11})$$

where

$$\lambda^a = g^{a\bar{b}} \frac{\partial}{\partial \lambda^{\bar{b}}}. \quad (\text{A12})$$

In this representation all of the operators, including Q , act covariantly on the functions Ψ . The covariance is explicit if we regard Ψ as an element of the exterior algebra over a basis of forms $\lambda^{\bar{a}}$. In this case $\pi \lambda^{\bar{a}} = -i\nabla \lambda^{\bar{a}}$, where ∇ is the covariant derivative, and $Q = i\bar{\partial}^\dagger$, where $\bar{\partial}$ is the antiholomorphic exterior derivative.

The Hamiltonian is obtained from Q by the relation $H = \{Q, Q^*\}$, which by the use of equation (A9) and the commutation relations gives

$$H = -\frac{1}{2}\nabla^2 + \frac{1}{2}R_{\bar{a}b} \lambda^{\bar{a}} \lambda^b \quad (\text{A13})$$

where $R_{\bar{a}b}$ is the Ricci tensor.

De Alfaro et al. [9] have pointed out that alternative orderings of the two operators in the supercharge (A9) would make the representation non-covariant. However, there do exist alternative expressions for the supercharge which have the same classical limit,

$$Q = \lambda^a \pi_a - i\lambda^a W_a(z) + \lambda^{\bar{a}} U_{\bar{a}}(z) \quad (\text{A14})$$

The condition $Q^2 = 0$ is satisfied if $\partial W = 0$ and $\bar{\partial}^\dagger U = 0$. The simplest way to satisfy these conditions would be to make W a total derivative,

$$W_a = i\xi \partial_a (R_c^b [x^c, p_b]) \quad (\text{A15})$$

which gives additional contributions to the Hamiltonian of order $\hbar^3$. Other contributions to the Hamiltonian can exist if the manifold has non-trivial Homology.

APPENDIX B: GEOMETRICAL FORMULAE

This appendix reviews some formulae used to reduce the Riemannian connection ∇ onto a submanifold $\mathcal{M}$. We construct a basis of 1-forms $(\mathbf{n}^I, \mathbf{e}^a)$, with dual basis $(\mathbf{n}_I, \mathbf{e}_a)$, made up to include the normal forms $\mathbf{n}^I$ and vectors $\mathbf{e}^a$ tangential to $\mathcal{M}$. The metric is then

$$\mathbf{g} = \sigma_{ab} \mathbf{e}^a \otimes \mathbf{e}^b + \eta_{IJ} \mathbf{n}^I \otimes \mathbf{n}^J \quad (\text{B1})$$

where σ is the metric on $\mathcal{M}$. Any vector $\mathbf{X}$ can be decomposed into a tangential and normal parts,

$$\mathbf{X}_{\parallel} = \mathbf{e}^a(\mathbf{X}) \mathbf{e}_a \quad (\text{B2})$$

$$\mathbf{X}_{\perp} = \mathbf{n}^I(\mathbf{X}) \mathbf{n}_I \quad (\text{B3})$$

The extrinsic curvature $\mathbf{k}^I$ and extrinsic torsion tensor $\mathbf{a}_I{}^J$ are defined by

$$k^I{}_{ab} = (\nabla_a \mathbf{n}^I)_b \quad (\text{B4})$$

$$a_I{}^J{}_a = (\nabla_a \mathbf{n}^J)_I \quad (\text{B5})$$

The extrinsic curvature measures the expansion of the normal forms and it is always symmetric in the surface indices. The extrinsic torsion measures changes in the normals along lines drawn in the surface. The extrinsic torsion is antisymmetric if the normals are of unit length.

We will define the tangential covariant derivative by its action on the basis vectors,

$$\parallel \nabla_{\mathbf{X}} \mathbf{e}_a = (\nabla_{X_{\parallel}} \mathbf{e}_a)_{\parallel}, \quad \parallel \nabla_{\mathbf{X}} \mathbf{n}_I = (\nabla_{X_{\parallel}} \mathbf{n}_I)_{\perp} \quad (\text{B6})$$

With this definition the derivative is a metric connection for both σ and η , with torsion related to $\mathbf{a}_I{}^J$. We make repeated use of the following expression for the derivative of a normal directed tensor in the body of the text,

$$\parallel \nabla_a \omega_{IJ} = \partial_a \omega_{IJ} + a_I{}^K{}_a \omega_{KJ} + a_J{}^K{}_a \omega_{IK} \quad (\text{B7})$$

The curvature tensor of $\parallel \nabla$ will be denoted by $\parallel R$.

The reduction of the curvature tensor is a standard procedure, and has been described in this notation in reference [17]. The most useful formula is Gauss equation,

$$R_{abcd} = \parallel R_{abcd} - \eta_{IJ} k^I{}_{ac} k^J{}_{bd} + \eta_{IJ} k^I{}_{ad} k^J{}_{bc} \quad (\text{B8})$$

In the case where $\mathcal{M}$ is a complex submanifold of a Kähler manifold, the trace k^I of the extrinsic curvature tensor vanishes [18].

APPENDIX C: ADIABATIC APPROXIMATION SCHEMES

This appendix shows how the hamiltonian operator and other operators can be reduced by using almost degenerate perturbation theory [20]. The method is equivalent to the adiabatic, or Born-Oppenheimer, approximation scheme for the energy eigenstates.

The Hamiltonian is first separated into an unperturbed part and a perturbation

$$H = H_0 + H_1. \quad (\text{C1})$$

The eigenstates of H_0 are then grouped into eigenspaces that cover a narrow range of eigenvalues. A projection operator P_0 projects onto the lowest of these eigenspaces and $P_1 = 1 - P_0$. The equation for the eigenstate with energy E is standard (see [20] for example),

$$H_R P_0 \Psi = E P_0 \Psi \quad (\text{C2})$$

where

$$H_R = E_0 + P_0 H_1 P_0 + P_0 H_1 P_1 (E - H_0 - H_1)^{-1} P_1 H_1 P_0 \quad (\text{C3})$$

The leading order terms give

$$H_R = E_0 + P_0 H_1 P_0 + P_0 H_1 P_1 (E_0 - H_0)^{-1} P_1 H_1 P_0 + \dots \quad (\text{C4})$$

For an adiabatic approximation scheme, we look for an expansion of the wave function in the form

$$\Psi(x) = \sum_n \psi(x^a) f_n(x^I, x^a) \quad (\text{C5})$$

where the wave function depends strongly on coordinates x^I and weakly on coordinates x^a . We take an unperturbed Hamiltonian H_0 which commutes with the x^a and define the functions f_n by

$$H_0 f_n = E_n(x^a) f_n. \quad (\text{C6})$$

These can be normalised with respect to a reduced product

$$\langle f, g \rangle_\perp = \int f^*(x^a, x^I) g(x^a, x^I) d\mu(x^I). \quad (\text{C7})$$

with measure $d\mu(x^I)$ chosen to make H_0 self-adjoint.

The projection operator P_0 is given by

$$P_0 \Psi = \psi_0(x^a) \quad (\text{C8})$$

Inserting complete sets of states into equation (C4) gives the equation for ψ_0 ,

$$H_R \psi_0 = E \psi_0 \quad (\text{C9})$$

where the reduced Hamiltonian operator has the perturbative expansion

$$H_R = E_0 + \langle H_1 \rangle_{00} + \sum_m \langle H_1 \rangle_{0m} (E_0 - E_m)^{-1} \langle H_1 \rangle_{m0} + \dots \quad (\text{C10})$$

The matrix elements

$$\langle H_1 \rangle_{mn} = \langle f_m, H_1 f_n \rangle_{\perp} \quad (\text{C11})$$

are operators of the reduced theory. They also act on the E_n .

Other operators are reduced in a similar way. Given an operator Q , we follow (C3) and define

$$Q_R = P_0 Q P_0 + P_0 Q P_1 (E - H)^{-1} P_1 H P_0. \quad (\text{C12})$$

In the adiabatic approximation,

$$Q_R = \langle Q \rangle_{00} + \sum_n \langle Q \rangle_{0n} (E_0 - E_n)^{-1} \langle H_1 \rangle_{n0} + \dots \quad (\text{C13})$$

is an operator that acts on the states ψ_0 .

The reduced operator satisfies

$$Q_R P_0 \Psi = P_0 Q \Psi \quad (\text{C14})$$

where Ψ is any energy eigenstate with eigenvalue E . This reduction is not generally a homomorphism of the operator algebras, i.e. $(QQ')_R \neq Q_R Q'_R$ in general, but if Q and Q' are symmetries of the Hamiltonian, then it follows from (C12) that

$$(QQ')_R = Q_R Q'_R \quad (\text{C15})$$

In particular, if Q is a supersymmetry operator with $H = \{Q, Q^*\}$, then

$$H_R = \{Q_R, Q_R^*\} \quad (\text{C16})$$

This simplifies the problem of reducing the Hamiltonian operator.

REFERENCES

- [1] S. Coleman *Aspects of Symmetry* (Cambridge University Press, Cambridge 1985).
- [2] G. W. Gibbons and N. S. Manton, *Nucl. Phys.* **B274** (1986) 183
- [3] J. P. Gauntlett, *Nucl. Phys.* **B400** (1993) 103
- [4] A. Sen, *Phys. Lett.* **329B** (1994) 217
- [5] N. Manton, *Phys. Lett.* **110B** (1982) 54
- [6] R. S. Ward, *Phys. Letts.* **158B** (1985) 424
- [7] B. DeWitt, *Phys. Rev.* **85** (1952) 635
- [8] E. Witten, *Nucl. Phys.* **B302**, 253 (1982).
- [9] V. De Alfaro, S. Fubini, G. Furlan and M. Roncadelli, *Nucl. Phys.* **B296** (1988) 402.
- [10] P. Maraner, *J. Phys. A* **28** (1995) 2939; *Ann. Phys.* **246** (1996) 325
- [11] H. Jensen and H. Koppe, *Ann. Phys.* **63** (1971) 586
- [12] W. J. Zakrzewski and A. M. Din, *Nucl. Phys.* **B253** (1985) 77; W. J. Zakrzewski and W. J. Stokoe, *Z. Phys.* **C34** (1987) 491
- [13] P. J. Ruback, *Commun. Math. Phys.* **116**, 645 (1988)
- [14] L. Alvarez-Gaume and D. Z. Freedman, *Ann. Phys.* **134** (1981) 85
- [15] B. Zumino, *Phys. Lett.* **87B** (1979) 203
- [16] L. Alvarez-Gaume and D. Z. Freedman, *Commun. Math. Phys.* **80** (1981) 443
- [17] J. Grant and I. G. Moss, *Phys. Rev.* **D56** (1997) 6284
- [18] H. Blaine Lawson Jr, *Lectures on Minimal Submanifolds*, (Publish or Perish inc, New York 1987)
- [19] A. L. Besse, *Einstein Manifolds*, (Springer Verlag, Berlin 1987).
- [20] J. J. Sakuri *Modern Quantum Mechanics* (Addison-Wesley, 1994)